



Read our COVID-19 research and news.

**PERSPECTIVE** | VIEWPOINT: COVID-19

## Reducing transmission of SARS-CoV-2

Kimberly A. Prather<sup>1</sup>, Chia C. Wang<sup>2,3</sup>, Robert T. Schooley<sup>4</sup>

[+ See all authors and affiliations](#)

*Science* 26 Jun 2020:  
Vol. 368, Issue 6498, pp. 1422-1424  
DOI: 10.1126/science.abc6197

**Article**

**Figures & Data**

**Info & Metrics**

**eLetters**

 **PDF**



Correctly fitted masks are an important tool to reduce airborne transmission of SARS-CoV-2, particularly in enclosed spaces, such as on this Moscow Metro train in Russia.

PHOTO: SERGEI FADEICHEV/TASS VIA GETTY IMAGES

Respiratory infections occur through the transmission of virus-containing droplets ( $>5$  to  $10\ \mu\text{m}$ ) and aerosols ( $\leq 5\ \mu\text{m}$ ) exhaled from infected individuals during breathing, speaking, coughing, and sneezing. Traditional respiratory disease control measures are designed to reduce transmission by droplets produced in the sneezes and coughs of infected individuals. However, a

We use cookies to improve your experience and enable functionality and security of this site. Data may be shared with third parties. Further detail is available in our Privacy Policy. By clicking "OK" you consent to our use of cookies and use of data. You can modify your cookie preferences by clicking "Manage Cookies".

OK

Manage  
Cookies

mission must be implemented, including universal masking and regular, widespread testing to identify and isolate infected asymptomatic individuals.

Humans produce respiratory droplets ranging from 0.1 to 1000  $\mu\text{m}$ . A competition between droplet size, inertia, gravity, and evaporation determines how far emitted droplets and aerosols will travel in air (4, 5). Larger respiratory droplets will undergo gravitational settling faster than they evaporate, contaminating surfaces and leading to contact transmission. Smaller droplets and aerosols will evaporate faster than they can settle, are buoyant, and thus can be affected by air currents, which can transport them over longer distances. Thus, there are two major respiratory virus transmission pathways: contact (direct or indirect between people and with contaminated surfaces) and airborne inhalation.

In addition to contributing to the extent of dispersal and mode of transmission, respiratory droplet size has been shown to affect the severity of disease. For example, influenza virus is more commonly contained in aerosols with sizes below 1  $\mu\text{m}$  (submicron), which lead to more severe infection (4). In the case of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), it is possible that submicron virus-containing aerosols are being transferred deep into the alveolar region of the lungs, where immune responses seem to be temporarily bypassed. SARS-CoV-2 has been shown to replicate three times faster than SARS-CoV-1 and thus can rapidly spread to the pharynx, from which it can be shed before the innate immune response becomes activated and produces symptoms (6). By the time symptoms occur, the patient has transmitted the virus without knowing.

#### **SIGN UP FOR THE SCIENCE eTOC**

Get the latest issue of *Science* delivered right to you!

Identifying infected individuals to curb SARS-CoV-2 transmission is more challenging compared to SARS and other respiratory viruses because infected individuals can be highly contagious for several days, peaking on or before symptoms occur (2, 7). These “silent shedders” could be critical drivers of the enhanced spread of SARS-CoV-2. In Wuhan, China, it has been estimated that undiagnosed cases of COVID-19 infection, who were presumably asymptomatic, were responsible for up to 79% of viral infections (3). Therefore, regular, widespread testing is essential to identify and isolate infected asymptomatic individuals.

Airborne transmission was determined to play a role during the SARS outbreak in 2003 (1, 4). However, many countries have not yet acknowledged airborne transmission as a possible pathway for SARS-CoV-2 (1). Recent studies have shown that in addition to droplets, SARS-CoV-2 may also be transmitted through aerosols. A study in hospitals in Wuhan, China, found

**Become a Member**

[Log In](#) [ScienceMag.org](#) 

to more than 100,000 droplets in emitted droplets per minute of speaking.

The Centers for Disease Control and Prevention (CDC) recommendations for social distancing of 6 feet and hand washing to reduce the spread of SARS-CoV-2 are based on studies of respiratory droplets carried out in the 1930s. These studies showed that large, ~100  $\mu\text{m}$  droplets produced in coughs and sneezes quickly underwent gravitational settling ([1](#)). However, when these studies were conducted, the technology did not exist for detecting submicron aerosols. As a comparison, calculations predict that in still air, a 100- $\mu\text{m}$  droplet will settle to the ground from 8 feet in 4.6 s, whereas a 1- $\mu\text{m}$  aerosol particle will take 12.4 hours ([4](#)). Measurements now show that intense coughs and sneezes that propel larger droplets more than 20 feet can also create thousands of aerosols that can travel even further ([1](#)). Increasing evidence for SARS-CoV-2 suggests the 6 feet CDC recommendation is likely not enough under many indoor conditions, where aerosols can remain airborne for hours, accumulate over time, and follow airflows over distances further than 6 feet ([5](#), [10](#)).


---

[Become a Member](#)

[Log In](#) [ScienceMag.org](#) 

# Masks reduce airborne transmission

Infectious aerosol particles can be released during breathing and speaking by asymptomatic infected individuals. No masking maximizes exposure, whereas universal masking results in the least exposure.



[Download high-res image](#) | [Open in new tab](#) | [Download Powerpoint](#)

[Become a Member](#)

[Log In](#) [ScienceMag.org](#) [Q](#)

☐ \PH' ☐ \LTC ☐ N/Sr ☐ 'E

In outdoor environments, numerous factors will determine the concentrations and distance traveled, and whether respiratory viruses remain infectious in aerosols. Breezes and winds often occur and can transport infectious droplets and aerosols long distances. Asymptomatic individuals who are speaking while exercising can release infectious aerosols that can be picked up by airstreams (10). Viral concentrations will be more rapidly diluted outdoors, but few studies have been carried out on outdoor transmission of SARS-CoV-2. Additionally, SARS-CoV-2 can be inactivated by ultraviolet radiation in sunlight, and it is likely sensitive to ambient temperature and relative humidity, as well as the presence of atmospheric aerosols that occur in highly polluted areas. Viruses can attach to other particles such as dust and pollution, which can modify the aerodynamic characteristics and increase dispersion. Moreover, people living in areas with higher concentrations of air pollution have been shown to have higher severity of COVID-19 (11). Because respiratory viruses can remain airborne for prolonged periods before being inhaled by a potential host, studies are needed to characterize the factors leading to loss of infectivity over time in a variety of outdoor environments over a range of conditions

Given how little is known about the production and airborne behavior of infectious respiratory droplets, it is difficult to define a safe distance for social distancing. Assuming SARS-CoV-2 virions are contained in submicron aerosols, as is the case for influenza virus, a good comparison is exhaled cigarette smoke, which also contains submicron particles and will likely follow comparable flows and dilution patterns. The distance from a smoker at which one smells cigarette smoke indicates the distance in those surroundings at which one could inhale infectious aerosols. In an enclosed room with asymptomatic individuals, infectious aerosol concentrations can increase over time. Overall, the probability of becoming infected indoors will depend on the total amount of SARS-CoV-2 inhaled. Ultimately, the amount of ventilation, number of people, how long one visits an indoor facility, and activities that affect airflow will all modulate viral transmission pathways and exposure (10). For these reasons, it is important to wear properly fitted masks indoors even when 6 feet apart. Airborne transmission could account, in part, for the high secondary transmission rates to medical staff, as well as major outbreaks in nursing facilities. The minimum dose of SARS-CoV-2 that leads to infection is unknown, but airborne transmission through aerosols has been documented for other respiratory viruses, including measles, SARS, and chickenpox (4).

Airborne spread from undiagnosed infections will continuously undermine the effectiveness of even the most vigorous testing, tracing, and social distancing programs. After evidence revealed that airborne transmission by asymptomatic individuals might be a key driver in the global spread of COVID-19, the CDC recommended the use of cloth face coverings. Masks provide a critical

[Become a Member](#)

[Log In](#) [ScienceMag.org](#) 

the other common viruses, such as health care settings, airplanes, restaurants, and other crowded places with reduced ventilation. The aerosol filtering efficiency of different materials, thicknesses, and layers used in properly fitted homemade masks was recently found to be similar to that of the medical masks that were tested (14). Thus, the option of universal masking is no longer held back by shortages.

From epidemiological data, places that have been most effective in reducing the spread of COVID-19 have implemented universal masking, including Taiwan, Japan, Hong Kong, Singapore, and South Korea. In the battle against COVID-19, Taiwan (population 24 million, first COVID-19 case 21 January 2020) did not implement a lockdown during the pandemic, yet maintained a low incidence of 441 cases and 7 deaths (as of 21 May 2020). By contrast, the state of New York (population ~20 million, first COVID case 1 March 2020), had a higher number of cases (353,000) and deaths (24,000). By quickly activating its epidemic response plan that was established after the SARS outbreak, the Taiwanese government enacted a set of proactive measures that successfully prevented the spread of SARS-CoV-2, including setting up a central epidemic command center in January, using technologies to detect and track infected patients and their close contacts, and perhaps most importantly, requesting people to wear masks in public places. The government also ensured the availability of medical masks by banning mask manufacturers from exporting them, implementing a system to ensure that every citizen could acquire masks at reasonable prices, and increasing the production of masks. In other countries, there have been widespread shortages of masks, resulting in most residents not having access to any form of medical mask (15). This striking difference in the availability and widespread adoption of wearing masks likely influenced the low number of COVID-19 cases.

Aerosol transmission of viruses must be acknowledged as a key factor leading to the spread of infectious respiratory diseases. Evidence suggests that SARS-CoV-2 is silently spreading in aerosols exhaled by highly contagious infected individuals with no symptoms. Owing to their smaller size, aerosols may lead to higher severity of COVID-19 because virus-containing aerosols penetrate more deeply into the lungs (10). It is essential that control measures be introduced to reduce aerosol transmission. A multidisciplinary approach is needed to address a wide range of factors that lead to the production and airborne transmission of respiratory viruses, including the minimum virus titer required to cause COVID-19; viral load emitted as a function of droplet size before, during, and after infection; viability of the virus indoors and outdoors; mechanisms of transmission; airborne concentrations; and spatial patterns. More studies of the filtering efficiency of different types of masks are also needed. COVID-19 has inspired research that is already leading to a better understanding of the importance of airborne transmission of respiratory disease.

---

[Become a Member](#)

[Log In](#) [ScienceMag.org](#) 



1. [↩](#) L. Morawska, J. Cao, *Environ. Int.* **139**, 105730 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)
2. [↩](#) E. L. Anderson *et al.*, *Risk Anal.* **40**, 902 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)
3. [↩](#) S. Asadi *et al.*, *Aerosol Sci. Technol.* **54**, 635 (2020). [CrossRef](#) [Google Scholar](#)
4. [↩](#) R. Tellier *et al.*, *BMC Infect. Dis.* **19**, 101 (2019). [CrossRef](#) [PubMed](#) [Google Scholar](#)
5. [↩](#) R. Mittal, R. Ni, J.-H. Seo, *J. Fluid Mech.* 10.1017/jfm.2020.330 (2020). [Google Scholar](#)
6. [↩](#) H. Chu *et al.*, *Clin. Infect. Dis.* 10.1093/cid/ciaa410 (2020). [Google Scholar](#)
7. [↩](#) X. He *et al.*, *Nat. Med.* **26**, 672 (2020). [PubMed](#) [Google Scholar](#)
8. [↩](#) Y. Liu *et al.*, *Nature* 10.1038/s41586-020-2271-3 (2020). [Google Scholar](#)
9. [↩](#) V. Stadnytskyi *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 202006874 (2020). [Google Scholar](#)
10. [↩](#) G. Buonanno, L. Stabile, L. Morawska, *Environ. Int.* **141**, 105794 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)
11. [↩](#) E. Conticini, B. Frediani, D. Caro, *Environ. Pollut.* **261**, 114465 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)
12. [↩](#) N. H. L. Leung *et al.*, *Nat. Med.* **26**, 676 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)
13. [↩](#) J. F.-W. Chan *et al.*, *Clin. Infect. Dis.* 10.1093/cid/ciaa644 (2020). [Google Scholar](#)
14. [↩](#) A. Konda *et al.*, *ACS Nano* 10.1021/acsnano.0c03252 (2020). [Google Scholar](#)
15. [↩](#) C. C. Leung, T. H. Lam, K. K. Cheng, *Lancet* **395**, 945 (2020). [CrossRef](#) [PubMed](#) [Google Scholar](#)

**Acknowledgments:** The authors thank S. Strathdee, D. Petras, and L. Marr for helpful discussions. K.A.P. is supported by the NSF Center for Aerosol Impacts on Chemistry of the Environment (CHE1801971). R.T.S. is supported by the National Institute of Allergy and Infectious Diseases (R01 AI131424). C.C.W. is supported by the Ministry of Science and Technology (MOST 108-2113-M-110-003) and the Higher Education Sprout Project of the Ministry of Education, Taiwan, ROC.

[View Abstract](#)

---

### Recommended articles from TrendMD

Airborne transmission of SARS-CoV-2  
Kimberly A. Prather *et al.*, *Science*

How should data on airborne transmission of  
SARS-CoV-2 change occupational health  
guidelines?

[Become a Member](#)

[Log In](#) [ScienceMag.org](#) [Q](#)

☐ Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS–coronavirus 2  
Jianzhong Shi et al., Science, 2020

Combination prevention for COVID-19  
Myron S. Cohen et al., Science, 2020

Consensus for prevention and management of coronavirus disease 2019 (COVID-19) for neurologists

Huijuan Jin et al., Stroke and Vascular Neurology, 2020

Flattening the curve in COVID-19 using personalised protective equipment: lessons from air pollution

Sanjay Rajagopalan et al., Heart, 2020

Breast feeding at the time of COVID-19: do not forget expressed mother's milk, please

Riccardo Davanzo, Arch Dis Child Fetal Neonatal Ed, 2020

Powered by **TREND MD**



## Science

Vol 368, Issue 6498  
26 June 2020

[Table of Contents](#)

[Print Table of Contents](#)

[Advertising \(PDF\)](#)

[Classified \(PDF\)](#)

[Masthead \(PDF\)](#)

## ARTICLE TOOLS

- ☐ Email
- ☐ Print
- ☐ Alerts
- ☐ Share

- ☐ Download Powerpoint
- ☐ Request Permissions
- ☐ Citation tools

## MY SAVED FOLDERS

- ☐ Save to my folders

## STAY CONNECTED TO SCIENCE

- [Facebook](#)
- [Twitter](#)

## RELATED CONTENT

Research Articles

**Become a Member**

[Log In](#) [ScienceMag.org](#)





## Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses

Editorials

### New coronavirus outbreak: Framing questions for pandemic prevention

Reports

### Antibody signature induced by SARS-CoV-2 spike protein immunogens in rabbits

Reports

### A rapidly deployable individualized system for augmenting ventilator capacity

Reports

### Using influenza surveillance networks to estimate state-specific prevalence of SARS-CoV-2 in the United States

#### RELATED INFORMATION

- [PODCAST Stopping the spread of COVID-19, and arctic adaptations in sled dogs](#)

#### SIMILAR ARTICLES IN:

- [PubMed](#)
- [Google Scholar](#)

#### CITING ARTICLES IN:

- [Web of Science \(37\)](#)

[Become a Member](#)

[Log In](#) [ScienceMag.org](#)



## Related Jobs

### Cancer Biology - Assistant Professor Tenure-Track University of Minnesota – Twin Cities

University of Minnesota  
Minneapolis, Minnesota

### Endowed Professor in Cancer Biology

University of Notre Dame  
Notre Dame

### Tenure-track Assistant Professor/Associate Professor

The University of Hong Kong  
Hong Kong (HK)

[MORE JOBS ▶](#)

### NAVIGATE THIS ARTICLE

- [Article](#)
  - [References and Notes](#)
- [Figures & Data](#)
- [Info & Metrics](#)

[Become a Member](#)

[Log In](#) [ScienceMag.org](#)

## Read the Latest Issue of *Science*

9 October 2020

Vol 370, Issue 6513



[Table of Contents](#)

### FEATURE

**Sweden's gamble**

### SPACE POLICY

**U.S. policy puts the safe development of space at risk**

### FICTION

**Plotting a path to a more sustainable future**

### SCI COMMUN

**News at a glance**

### COMMUNITY ECOLOGY

**Novel communities are a risky business**

### WORKING LIFE

**A pandemic journey**

## Get Our E-Alerts

Receive emails from *Science*. [See full list](#)

- ☒ *Science* Table of Contents
- ☒ *Science* Daily News
- ☒ Weekly News Roundup
- ☒ *Science* Editor's Choice
- ☒ First Release Notification
- ☒ *Science* Careers Job Seeker

Italy



Email address\*

- ☐ I also wish to receive emails from AAAS/*Science* and *Science* advertisers, including information on

[Become a Member](#)

[Log In](#) [ScienceMag.org](#)

**Sign up today**

---

Required fields are indicated by an asterisk (\*)

**About Us**

- Journals
- News from Science
- Leadership
- Team Members
- Work at AAAS

**For Advertisers**

- Advertising Kits
- Awards and Prizes
- Custom Publishing
- Webinars

**For Authors**

- Submit
- Information for Authors
- Editorial Policies

**For Librarians**

- Manage Your Institutional Subscription
- Information for Librarians
- Request a Quote
- FAQs

**Related Sites**

- AAAS.org
- EurekAlert!
- Science in the Classroom
- Science Magazine Japanese

**Help**

- Access and Subscriptions
- Order a Single Issue
- Reprints and Permissions
- Contact Us
- Accessibility



**Become a Member**

Log In ScienceMag.org

- Terms of Service
- Privacy Policy
- Contact AAAS